Recombinant Protein A-Cys, His

Catalog #: 1001-02

Lot #: See label



Description

Molecular Weight:

Approximately 34.8 kDa with $6\times$ His-tag and a Cys on C-terminus, a single non-glycosylated polypeptide chain containing 307 amino acids.

Source:

Escherichia coli.

Quantity:

10mg/100mg/1g

Protein per Gram Powder:*

See Label.

Concentration for 1 unity OD at 280 nm:

4.589 mg/ml

Specification

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Purity:

> 96 % by SDS-PAGE and RP-HPLC analyses.

Specificity:

The recombinant Protein A is a genetically engineered protein containing 5 IgG-binding regions of protein A. Cell wall binding region, cell membrane binding region and albumin binding region have been removed from the recombinant Protein A to ensure the maximum specific IgG binding. The recombinant Protein A is ideal for

purification of polyclonal or monoclonal IgG antibodies. Protein A binds to various human, mouse and rat IgG subclasses (e.g., human IgG1, IgG2, IgG4; mouse IgG2a, IgG2b, IgG3; rat IgG2c). It also binds to total IgG from rabbit, guinea pig, pig, dog and cat.

Endotoxin Level:

Less than 0.1 EU/ μg of Protein A-Cys, His as determined by LAL method.

Formulation:

Lyophilized from additive free solution.

Usage Guide

Stability & Storage:

- Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 3 months, -20 to -70 °C under sterile conditions after reconstitution.

Reconstitution:

Dissolve in distilled water or saline.

Shipping:


The product is shipped at ambient temperature. Upon receipt, store it immediately at -20 $^{\circ}$ C.

*Masses of the protein and powder are not equal.


Background

Protein A is a cell wall component produced by several strains of Staphylococcus aureus that consists of a single polypeptide chain and contains little or no carbohydrate. Recombinant Protein A is produced in E.Coli and functions essentially the same as native Protein A. It consists of 5 IgG-binding domains E-D-A-B-C aligned in series.

SDS-PAGE



RP-HPLC



Rev. 09/19/2023 V.3