

# Recombinant Human 4-1BB Receptor (rHu4-1BB Receptor)

## PrimeGene Technical Data Sheet

---

<b>Catalog Number:</b>	103-14R
<b>Source:</b>	<i>Escherichia coli</i> .
<b>Molecular Weight:</b>	Approximately 17.7 kDa, a single non-glycosylated polypeptide chain containing 166 amino acids.
<b>Quantity:</b>	5µg/20µg/1000µg
<b>AA Sequence:</b>	ERTRSLQDPC SNCPAGTFCD NNRNQICSPC PPNSFSSAGG QRTCDICRQC KGVFRTRKEC SSTSNAECDP TPGFHCLGAG CSMCEQDCKQ GQELTKKGCK DCCFGTFNDQ KRGICRPWTN CSLDGKSVLV NGTKERDVVC GPSPADLSPG ASSVTTPAPA REPGHS
<b>Purity:</b>	> 97 % by SDS-PAGE and HPLC analyses.
<b>Biological Activity:</b>	Fully biologically active when compared to standard. The biological activity is determined by its inhibitory effect of IL-8 production using human peripheral blood mononuclear cells. About 90 % of inhibition was seen using a concentration of 1 µg for both 4-1BB Ligand and 4-1BB Receptor.
<b>Physical Appearance:</b>	Sterile Filtered White lyophilized (freeze-dried) powder.
<b>Formulation:</b>	Lyophilized from a 0.2 µm filtered concentrated solution in 10 mM PB, pH 8.0, 150 mM NaCl.
<b>Endotoxin:</b>	Less than 1 EU/µg of rHu4-1BB Receptor as determined by LAL method.
<b>Reconstitution:</b>	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
<b>Shipping:</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage:</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"><li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li><li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li><li>● 3 months, -20 to -70 °C under sterile conditions after reconstitution.</li></ul>
<b>Usage:</b>	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. <b>NOT FOR HUMAN USE.</b>

---

### ***Human 4-1BB Receptor***

4-1BB receptor, also named TNFRSF9 is a member of the TNF superfamily of receptors. It is mainly expressed on the surface of a variety of T cells, but also found in B cells, monocytes, and various transformed cell lines. 4-1BB receptor binds to 4-1BBL, and they co-stimulate activity for activated T cells. Signaling by 4-1BB Receptor has been implicated in the antigen-presentation process and generation of cytotoxic T cells. Crosslinking of 4-1BB Receptor enhances T cell proliferation, IL-2 secretion survival and cytolytic activity. Further, it can enhance immune activity to eliminate tumors in mice.