

PrimeGene Technical Data Sheet

Catalog Number:	103-14
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 19.4 kDa, a single non-glycosylated polypeptide chain containing 184 amino acids.
Quantity:	5µg/20µg/1000µg
AA Sequence:	REGPELSPDD PAGLLDLRQG MFAQLVAQNV LLIDGPLSWY SDPGLAGVSL TGGLSYKEDT KELVVAKAGV YYVFFQLELR RVVAGEGSGS VSLALHLQPL RSAAGAAALA LTVDLPPASS EARNSAFGFQ GRLLHLSAGQ RLGVHLHTEA RARHAWQLTQ GATVLGLFRV TPEIPAGLPS PRSE
Purity:	> 95 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Fully biologically active when compared to standard. The ED ₅₀ as determined by stimulation of IL-8 production using human PBMC is less than 10 ng/ml, corresponding to a specific activity of > 1.0 × 10 ⁵ IU/mg.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4.
Endotoxin:	Less than 1 EU/µg of rHu4-1BBL as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE.

Human 4-1BB Ligand

4-1BBL is a member of the tumor necrosis factor (TNF) receptor family. This receptor contributes to the clonal expansion, survival, and development of T cells. In addition, 4-1BBL expression is found on dendritic cells, follicular dendritic cells, natural killer cells, granulocytes and cells of blood vessel walls at sites of inflammation. CD137 has been shown to interact with TRAF2. The human 4-1BBL gene codes for a 254 amino acid type II transmembrane containing a 28 amino acid cytoplasmic domain, a 21 amino acid transmembrane domain, and a 205 amino acid extracellular domain (ECD). The human 4-1BBL ECD shares 32 % and 35 % a.a. identity with murine and rat ECD.