

Recombinant Human Cysteine-rich Angiogenic Inducer 61 (rHuCYR61)

PrimeGene Technical Data Sheet

Catalog Number:	105-18
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 39.4 kDa, a single non-glycosylated polypeptide chain containing 357 amino acids.
Quantity:	5µg/20µg/1000µg
AA Sequence:	TCPAACHCPL EAPKCAPGVG LVRDGC GCK VCAKQLNEDC SKTQPCDHTK GLECNFGASS TALKGICRAQ SEGRPCEYNS RIYQNGESFQ PNCKHQCTCI DGAVGCIPLC PQELSLPNLG CPNRLVKVT GQCCEEWVCD EDSIKDPMED QDGLLGKELG FDASEVELTR NNELIAVGKG SSLKRLPVFG MEPRILYNPL QGQKCIQTT SWSQCSKTCG TGISTRVTND NPECRLVKET RICEVRPCGQ PVYSSLKKGK KCSKTKKSPE PVRFTYAGCL SVKKYRPKYC GSCVDGRCT PQLTRTVKMR FRCEGETFS KNVMMIQSCK CNYNCPHANE AAFPFYRLFN DIHKFRD
Purity:	> 95 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Fully biologically active when compared to standard. The ED ₅₀ as determined by a cell proliferation assay using murine Balb/3T3 cells is less than 3.0 µg/ml, corresponding to a specific activity of > 330 IU/mg.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in citrate buffer solution, 300 mM NaCl, pH 3.0.
Endotoxin:	Less than 1 EU/µg of rHuCYR61 as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE.

Human Cysteine-rich Angiogenic Inducer 61

Cysteine-rich angiogenic inducer 61 (Cyr61) encoded by the Cyr61 gene is a dynamically expressed, multifunctional matricellular protein and it is also a secreted, extracellular matrix (ECM)-associated signaling protein of the CCN family. Cyr61 plays essential roles in cardiovascular development during embryogenesis and regulates inflammation, wound healing and fibrogenesis in the adult. Aberrant CCN1 expression is associated with myriad pathologies, including various cancers and diseases associated with chronic inflammation. Mature human Cyr61 shares 93 % amino acid sequence identity with mouse and rat Cyr61. Cyr61 consists of four domains. There are an IGFBP domain, a VWF type C domain, a TSP type I domain, and a cysteine knot domain.