

PrimeGene Technical Data Sheet

Catalog Number:	141-01A
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 17.8 kDa, a single non-glycosylated polypeptide chain containing 156 amino acids.
Quantity:	2 μ g/10 μ g/1000 μ g
AA Sequence:	SAPHSFQNNL RYKLIRIVKQ EFIMNDSLQ NIYVDMDRIH LKAASLNDLQ LEVKFDMYAY SSGGDDSKYP VTLKVSNTQL FVSAQGEDKP VLLKEIPETP KLITGSETDL IFFWEKINSK NYFTSAAFPE LLATKEQSQ VHLARGLPSM IDFQIS
Purity:	> 97 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Fully biologically active when compared to standard. The ED ₅₀ as determined by a cell proliferation assay using murine D10S cells is less than 0.005 ng/ml, corresponding to a specific activity of > 2.0 \times 10 ⁸ IU/mg.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 μ m filtered concentrated solution in 20 mM Tris, 150 mM NaCl, pH 8.5.
Endotoxin:	Less than 1 EU/ μ g of rRtIL-1 α as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at \leq -20 °C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE.

Rat Interleukin-1 alpha

Interleukin-1 alpha (IL-1 α) is a non-secreted proinflammatory cytokine produced mainly by activated macrophages, as well as neutrophils, epithelial cells, and endothelial cells. It possesses metabolic, physiological, haematopoietic activities, and plays one of the central roles in the regulation of the immune responses. Both IL-1 α and IL-1 β binds to the same receptor and has similar but not identical biological properties. Recombinant rat IL-1 alpha contains 150 amino acids which is a single non-glycosylated polypeptide.