

PrimeGene Technical Data Sheet

Catalog Number:	147-07
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 16.6 kDa, a single non-glycosylated polypeptide chain containing 141 amino acids.
Quantity:	2μg/10μg/1000μg
AA Sequence:	SESLVVCDVA EDLVEKLRKF RFRKETHNAA IIMKIDKDKR LVVLDEELEG VSPDELKDEL PERQPRFIVY SYKYQHDDGR VSYPLCFIFS SPLGCKPEQQ MMYAGSKNKL VQTAELTKVF EIRNTEDLTE EWLREKLGFF H
Purity:	> 98 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Data is not available.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 μm filtered concentrated solution in PBS, pH 7.4.
Endotoxin:	Less than 1 EU/μg of rRtGMF-β as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE.

Rat Glia Maturation Factor beta

The glia maturation factor beta belongs to the actin-binding proteins ADF family, GMF subfamily. It contains an ADF-H domain, but the research of crystallography and NMR reveals that there are structures different between human and mouse ADF-H domain. GMF-β is involved in the differentiation, maintenance, and regeneration of the nervous system. It also inhibition of proliferation of tumor cells.