

## PrimeGene Technical Data Sheet

---

|                                 |  |
|---------------------------------|--|
| <b>Catalog Number:</b>          | 301-01G  |
| <b>Source:</b>                  | <i>Escherichia coli</i> .  |
| <b>Molecular Weight:</b>        | Approximately 22 kDa, a single non-glycosylated polypeptide chain containing 191 amino acids.  |
| <b>Quantity:</b>                | 100µg/500µg/1mg  |
| <b>AA Sequence:</b>             | FPTIPLSRLF DNAMLRAHRL HQLAFDITYQE FEEAYIPKEQ KYSFLQNPQT SLCFSESIPT<br>PSNREETQOK SNLELLRISL LLIQSWLEPV QFLRSVFANS LVIYASDSNV YDLLKDLEEG<br>IQTLMGRLED GSPRTGQIFK QTYSKFDTNS HNDDALLKNY GLLYCFRDKM DKVETFLRIV<br>QCRSVEGSCG F   |
| <b>Purity:</b>                  | > 96 % by SDS-PAGE and HPLC analyses.  |
| <b>Biological Activity:</b>     | Fully biologically active when compared to standard. The ED <sub>50</sub> as determined by a cell proliferation assay using rat Nb2-11 lymphoma cells is less than 0.1 ng/ml, corresponding to a specific activity of > 1.0 × 10 <sup>7</sup> IU/mg.   |
| <b>Physical Appearance:</b>     | Sterile Filtered White lyophilized (freeze-dried) powder.  |
| <b>Formulation:</b>             | Lyophilized from a 0.2 µm filtered solution in PBS, pH 7.0, containing 3 % Mannitol, 0.02 % Tween-20.  |
| <b>Endotoxin:</b>               | Less than 0.1 EU/µg of rHuGH as determined by LAL method.  |
| <b>Reconstitution:</b>          | We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions. |
| <b>Shipping:</b>                | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.  |
| <b>Stability &amp; Storage:</b> | <b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"><li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li><li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li><li>● 3 months, -20 to -70 °C under sterile conditions after reconstitution.</li></ul>                                 |
| <b>Usage:</b>                   | This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. <b>NOT FOR HUMAN USE.</b>   |

---

### ***Human Growth Hormone***

Growth hormone (GH), also known as somatotropin, is a member of a family of growth factors that includes prolactin, placental lactogens, proliferins, and somatolactin. It is a peptide hormone that stimulates growth, cell reproduction and regeneration in humans and other animals. Growth hormone is a 191-amino acid, single-chain polypeptide that is synthesized, stored, and secreted by somatotropic cells within the lateral wings of the anterior pituitary gland. The pulsatile release of GH into circulation is regulated by the concerted actions of the hypothalamic hormones-GH-releasing hormone (GHRH) and somatostatin (SST) as well as by signals from the periphery-ghrelin and leptin. The human GH cDNA encodes a 217 amino acid (a.a.) residue precursor protein with a 26 a.a. putative signal peptide. By alternative splicing, at least four isoforms of GH have been identified.