

PrimeGene Technical Data Sheet

Catalog Number:	601-35
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 36.0 kDa, a single non-glycosylated polypeptide chain containing 330 amino acids, including the 317 residues of full-length Sox2 and a 13-residue C-terminal TAT peptide (GGYGRKKRRQRRR).
Quantity:	5µg/25µg/1000µg
AA Sequence:	MYNMMETELK PPGPQQTSGG GGGNSTAAAA GGNQKNSPDR VKRPMNAFMV WSRGQRRKMA QENPKMHNSE ISKRLGAEWK LLETETKRPF IDEAKRLRAL HMKEHPDYKY RPRRKTTLML KKDKYTLPGG LLAPGGNSMA SGVGVGAGLG AGVNQRMSY AHMNGWSNGS YSMMQDQLGY PQHPGLNAHG AAQMMPMHRY DVSALQYNSM TSSQTYMNGS PTYSMSYSQQ GTPGMALGSM GSVVKSEASS SPPVVTSSSH SRAPCQAGDL RDMISMYPG AEVPEAAPS RLHMSQHYQS GPVPGTAING TLPLSHMGY GRKKRRQRRR
Purity:	> 95 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Data Not Available.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in 2 × PBS, pH 7.4, with 5% Trehalose.
Endotoxin:	Less than 1 EU/µg of rHuSOX2-TAT as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/ml. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● 12 months from date of receipt, -20 to -70 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE.

Human SOX2-TAT

Sox2, also named SRY (sex determining region Y)-box 2, is belonging to the Sox family and it is encoded by the Sox gene in human. This protein family shares highly conserved DNA binding domains known as HMG (High-mobility group) box domains containing approximately 80 amino acids. Sox2 plays a role in maintenance of embryonic and neural stem cells and holds great promise in research involving induced pluripotency, an emerging and very promising field of regenerative medicine. Mature human Sox2 shares 100 amino acid sequence identity with murine and rat Sox2. Recombinant human Sox2-TAT expressed in *E. coli* is a 36 kDa protein containing 330 amino-acid residues, including the 317 residues of full-length Sox2 and a 13-residue C-terminal TAT peptide (GGYGRKKRRQRRR).