

PrimeGene Technical Data Sheet

Catalog Number:	602-19
Source:	<i>Escherichia coli</i> .
Molecular Weight:	Approximately 28.1 kDa, a single non-glycosylated polypeptide chain containing 243 amino acids.
Quantity:	20µg/100µg/1000µg
AA Sequence:	DEPPQSPWDR VKDLATVYVD VLKDSGRDYV SQFEGSALGK QLNKLLDNW DSVTSTFSKL REQLGPVTQE FWDNLEKETE GLRQEMSKDL EEVKAKVQPY LDDFQKKWQE EMELYRQKVE PLRAELQEGA RQKLHELQEK LSPLGEEMRD RARAHVDALR THLAPYSDEL RQRLAARLEA LKENGGARLA EYHAKATEHL STLSEKAKPA LEDLRQGLLP VLESFKVSFL SALEEYTKKL NTQ
Purity:	> 98 % by SDS-PAGE and HPLC analyses.
Biological Activity:	Data Not Available.
Physical Appearance:	Sterile Filtered White lyophilized (freeze-dried) powder.
Formulation:	Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4.
Endotoxin:	Less than 0.1 EU/µg of rHuApoA-I as determined by LAL method.
Reconstitution:	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage:	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none">● Refer to lot specific COA for the Use by Date when stored at ≤ -20 °C as supplied.● 1 month, 2 to 8 °C under sterile conditions after reconstitution.● 3 months, -20 to -70 °C under sterile conditions after reconstitution.
Usage:	This material is offered by Shanghai PrimeGene Bio-Tech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE.

Human Apolipoprotein A-I

Apolipoprotein A-I encoded by the APOA1 gene is the major protein component of high density lipoprotein (HDL) in plasma. It has a specific role in lipid metabolism. Apolipoprotein A-I participates in the reverse transport of cholesterol from tissues to the liver for excretion by promoting cholesterol efflux from tissues and by acting as a cofactor for the lecithin cholesterol acyltransferase (LCAT). Apolipoprotein A-I is also isolated as a prostacyclin (PGI₂) stabilizing factor and thus may have an anticlotting effect. Apolipoprotein A-I has been shown to interact with ABCA1, GPLD1 and PLTP. Apolipoprotein A-I binds to lipopolysaccharide or endotoxin and has a major role in the anti-endotoxin function of HDL. Mature human ApoA-1 shares 65 % and 62 % a.a. sequence identity with murine and rat ApoA-1, respectively. It has an approximate molecular weight of 28.1 kDa.